
Shewanella algae meningitis and early onset neonatal sepsis : A case report of Libyan newborn And literature review

Fathia. A .Almurabet ,Kheria.O.Almamori

Neonatal intensive care unit , Pediatric department, Zawia Teaching hospital, faculty of Medicine, Zawia university –Libya

correspondence ;Murabetf@ Gmail.com

Abstract

Neonatal sepsis still a main leading cause of morbidity and mortality among neonates, especially in the developing countries ,Most cases of Neonatal sepsis in neonatal intensive care unit are attributed to GB streptococcus, klebsiella pneumonia and other members of Enterobacteriaceae family. Shewanella algae (S.algae) is a gram –negative saprophytic bacillus, commonly associated with the marine environment ,which has been isolated from humans. Early onset neonatal sepsis caused by S.algae is uncommon. We report a first case of Shewanella. Algae meningitis in newborn with early onset neonatal sepsis which was responding to antimicrobial therapy and associated with good early out come and no late neurodevelopmental sequel.

Key words: shewanella algae, neonatal meningitis, Early onset neonatal sepsis . Cerebrospinal fluid culture .blood culture , good outcome .

Background

Neonatal sepsis, a blood stream infection occurring during the first month of life ,is associated with increased mortality and morbidity .¹Early onset neonatal sepsis (EONS)develop within three days of birth .The organisms associated with it are derived from the mother during the intrapartum period ;these are Group B

streptococci klebsiella, Escherichia coli, streptococcus viridians ,enterococci,Staphylococcus aureus, Pseudomonas aeruginosa and the other gram negative bacilli .^{2,3}shewanella spp ;saprophytic non fermenting gram –negative bacilli ,are an uncommon cause of human disease associated with three characteristic patterns of human infection ;i.e., soft tissue infection, bacteremia and neonatal sepsis

^{4,6,12}Here ,We report a case of neonatal meningitis and early onset neonatal sepsis caused by shewanella algae.

What this study adds:

1.what is known about this subject ?

Shewanella algae is a marine bacteria and an uncommon cause of infection in humans caused by exposure to sea water .However, these infections are often associated with high mortality .Prematurity is a known risk factor.

2.what new information is offered in this report ?

We report an uncommon case of meningitis with early onset neonatal sepsis caused by shewanella algae in a newborn delivered by caesarean section at 34 weeks of gestation.

S. algae is saprophytic bacterium and a novel pathogen of neonatal sepsis and very rare cause of neonatal meningitis particularly in an institutional setup ,where exposure to contaminated water is unlikely .

3 .what are the implications for research ,policy ,or practice. ?

This case highlights the importance of a clinical suspicion and microbiology laboratory in indentifying the correct

etiological agent for neonatal meningitis and sepsis (Shewanella algae in our case) ,therefore facilitating the use of an appropriate antibiotics, which resulted in a better and satisfactory clinical early and late out come .

Case details:

A39 years old para four diabetic lady with previous four caesarean sections presented at 34 weeks gestation in labor ,she has history of intrauterine fetal death at 36 weeks and history of abortion at 18 weeks. no history of premature rupture of membrane .Emergency lower segment caesarean section was done and 3.235kg female baby was delivered. Apgar score was 5/10 at one minute and 9/10 at 5 minute then the baby was transferred to the special care baby unit for management as infant of diabetic mother where the baby develop sign of mild respiratory distress (Transit tachypnea of newborn TTN). Oxygen supply with oxyhood , intravenous fluid was started . Complete blood count revealed WBC 24 Hgb18.7g/dl platelets 266 Lymphocyte 25.8% Neutrophils 62.8% blood sugar, Serum Calcium monitoring was done and managed accordingly, Early enteral (trophic) feeding was started by Nasogastric tube. On the second day

respiratory distress was subsided (Transit tachypnea of newborn).oral feeding by sucking was started ,she developed hypocalcaemia and hypomagnacemia . treatment was given ,At the end of 2nd day of life she was well ,jaundice, serum bilirubin was at phototherapy level , C reactive protein (CRP) was negative .On the 3th day, she was unwell depressed activity, hypoxic, O₂saturation by pulse oximetry 70%, Febrile 39.5 °C tachypnic RR 65/min, tachycardia 170 beat /min systemic examination was normal .the baby was transferred to the intensive care unit where CBC ,CRP Blood sugar , S.Calcium ,B .Urea and electrolyte .full septic screen was done (Lumber puncture ,suprapubic aspiration for 2 urine culture and sensitivity, blood for culture and sensitivity ,chest X ray),parenteral antibiotic Ampicilin 100mg/kg day divided 12 hourly and gentamycin 5mg/kg /day iv divided 12 hourly was started .investigations results revealed total white cell count $21.1 \times 10^3/\mu\text{l}$, red blood cells $5.05 \times 10^6/\mu\text{l}$, hemoglobin 17.1 g /dL ,platelet $173 \times 10^3 /\mu\text{l}$.the differential leucocytes count (Neutrophils 82.4%,lymphocyte 13.8%,mixed 3.8%.immature Neutrophils $17.4 \times 10^3/\mu\text{l}$.CRP was raised 4.8mg /dl .Ampicilin and gentamycin was started as

empirical therapy .blood sugar at time of Lumber puncture was 44 mg/dl, cerebrospinal fluid sugar was <10mg/dl, cerebrospinal fluid protein was >300mg/dl .the baby was stable Ampicilin dose was increased to 150mg /kg 12 hourly intravenously .After 24 hours the baby general condition was unsatisfactory ,lethargic ,mottled skin ,developed prolonged apnea stopped by mechanical stimulation. complete blood count , CRP , blood sugar ,urea ,electrolyte s, serum calcium , blood culture was taken . Ampicilin and gentamycin were discontinued. cefotaxime (claforan)and Amikacin were started . CRP was 2.4mg /dl . WBC $20.1 \times 10^3/\mu\text{L}$ neutrophils % 65% lymphocyte%25.9% HGB 16 gm/dl S.Calcium was 7.7 mg/dl, blood sugar was 188mg/dl blood urea 25mg/dL, sodium 146 mmol/l .potassium 4.2mmol/l .treatment of hypocalcaemia was done. Blood and cerebrospinal fluid cultures were done by automated bioMerieux customer and VITEK 2- compact and gave a positive result after 24 hours where Gram negative bacilli which produced β – haemolytic colonies with a tan pigment on blood agar media were isolated from CSF culture (figure1) which is growth of shewanella

algae. blood culture also revealed the growth of shewanella algae after 3 days.



Figure1:showing β haemolytic colonies with tan pigment of gram negative bacilli on blood agar media (shewanella algae)

The anti microbial susceptibility test reported that isolate was resistant to Ampicillin ,gentamycin ,Amoxicillin Clavulanic acid, and sensitive to cefotaxime ,Ceftazidime ,ceftriaxone.meropenem ,imipenem, ciprofloxacin . urine culture showed no growth .the patient became better, his vital signs and oxygen saturation

by pulse oximetry were normal. Neurological and systemic examination was normal, repeated blood sugar and serum calcium were normal when the cultures revealed growth of Shewanella algae we ask the mother about exposure to sea water and she reported that her house is too near to the sea and gave history of swimming in sea water frequently before delivery .Head ultrasound examination was normal .CRP after 3 days of starting cefotaxime(claforan)and amikacin was 1.2 mg/dl .one week after it was negative. (Two weeks of treatment was completed) the baby was well, accepting oral feeding. normal examination CRP negative . No growth in second blood culture. antibiotic stopped and discharged with good condition .on follow up as outpatient she was doing well .normal development. Now she is 5 years old with normal physical and neurological development.

Discussion:

The genus shewanella is an oxidative ,H₂S producing gram negative bacillus^{1, 4,6,7}_the two bio types ,S.algae and S.putrifaciens^{7,9} .S .putrefaciens are differentiated based on haemolytic property on blood agar, ability to ferment carbohydrates ,growth at 42 °C ,ability to grow on salmonella_ shigella agar

and 6.5% NaCl sodium chloride.^{1,4,9,10} shewanella spp are often isolated from natural habitats like water and soil. Human infection caused by this group of organism is rare .common infection caused by this organism include skin and soft tissue infection^{8,9,10,11,12,14,16} ,hepatobiliary disease^{6,13} ,and bacteremia^{4,9,13,18} , including neonatal sepsis,^{2,,5,10,11,12,14,15,16} .commonly ,infections caused by this organism are associated with exposure to environmental sources like water

The first cases of *S. algae* bacteremia were reported in Denmark (1999) in two patients with chronic leg ulcers. They had been exposed to the same marine environment .⁸Iwata etal reported a primary shewanella algae septicemia in a Japanese women undergoing hemodialysis¹⁸ . *S.algae* infections have been reported in children ,but there have been no previous reports of bacteremia .Ear infections are common ,and a series of 65 cases described children who have otitis media that occur during summer^{1,16} .^{*}Sardelic etal reported a case of ventriculitis and peritonitis caused by *S.algae* it was in Croatia in child with ventriculoperitoneal shunt¹⁷ ,The infection developed after contact with seawater and began as otitis¹⁴ . Tan CK etal reported a case of purulent pericarditis with greenish

pericardial effusion caused by shewanella algae.¹⁹

Shewanella algae bacteremia and associated cellulites in a patient with multiple myeloma was seen in Budap reported by Krsnik.²⁰

A case of bacteremia and suppurative vertebral osteomyelitis /discitis due to shewanella algae occurring after raw-fish consumption in Kansenshogakau Zasshi reported by Shimizu and Matsumura²¹ .

An outbreak of shewanella algae and shewanella putrefaciens infections caused by a shared measuring cup in general surgery unit in Korea was happen reported by OhHs,Kum etal²² .

Since traditional new born care practice like bathing or prelacteal feeding are wide in rural populations ,adherence to the standard obstetric care and health education would be imperative .In preventing this infections ,especially in the case of non institutional deliveries .in hospital setups, vigilant nursing care along with surveillance of hospital environment and facility for rapid laboratory diagnosis is likely to reduce the risk of shewanella infections¹⁵ ,in our patient the source of infection remained undetected .Although the infection was acquired postnatally ,water was unlikely to be a

source in this case as the baby was not exposed for bathing or pre lacteal feeding .More ever the enviromental screening of water sources in neonatal intensive care unit and delivery suite failed to grow shewanella spp. So ,the infection was most likely maternal acquired as the mother reported swimming in sea water before delivery , In Carolina, García –Fragoso etal reported S .algae bacteremia in extreme preterm baby with maternal history of swimming in sea water early in pregnancy.¹⁶ Shewanella bacteremia has been reported in infants premature ,and term neonates^{7,13,15,,16} ,Brink etal studied bacteremia in 24 patients , of which the majority were premature neonate (66% ,16 out of 24) ,^{2,7,,10,15,16}.Pnemonia and Respiratory distress were strongly associated with shewanella bacteremia ¹⁶,how ever in our case the baby has no respiratory illness apart from TTN in the first hours of life as in Charles etal reported a case of neonatal sepsis caused by S.algae has no associated respiratory diseases¹⁵. In the case series of infections *by* shewanella spp. However ,all the patients were adult and mainly presented with wound and soft infections which were chronic in nature^{9,18,19}.In all the previously reported cases of shewanella infection in adults ,children and neonates no case of meningitis

had been documented in the literature . Although S.algae has been isolated from various human samples¹ .Early onset neonatal sepsis[EONS] caused by this organism is un common. Around 13-28% mortality has been reported with shewanella infection .^{1,10} In spite , There is treatment failure due to emergence of resistance of carbapenem during therapy for shewanella algae bacteremia reported by Kim and his etal²³. Susceptibility to gentamycin ,cephalosporine and ciprofloxacin was seen in various studies.^{11,15,16} This is in accordance with our finding since in our isolated strain exhibited resistance to the Ampicilin ,gentamycin and highly sensitive to Amikacin ,cephalosporines and carbapenemes. The patient was improved clinically as we were starting gentamycin as empirical therapy and changed to Amikacin and cephalosporine in from of cefotaxime (claforan)when the patient became febrile and CRP was raising .and repeated blood culture result revealed no growth and finally the patient did not develop any early or late sequels which support the sensitivity of our isolated strain to these antibiotic treatment .So, early detection and initiation of proper treatment Of shewanella algae meningitis and sepsis results to successful outcome.

Conclusion

Shewanella algae is an unusual human pathogen. neonatal sepsis caused by shewanella algae is uncommon and neonatal meningitis caused by shewanella algae is very rare and was not reported in the literature before. Early diagnosis and proper management of neonatal meningitis caused by S.algae is associated with early good outcome and normal neurodevelopment.

References:

- 1.Holt HM ,Gahrn-Hansen B, Brun B.Shewanella putrefaciens :clinical and microbiological characteristics.Clin Microbiol Infect.2005;11:347-352.
- 2.Hornik CP,FortP,Clark RH, Watt K,Benjamin DK Jr,SmithPB.et al. Early and late onset sepsis in very- low birth weight infants from a large group of neonatal intensive care units. Early Hum Dev.2012;88(2):S69-74.[PMC Free article][Pub Med][Google Scholar]
- 3.BakerCJ,Barrett FF.Group B streptococcal infections in infants. The importance of the various serotypes.JAMA.1974;230:1158-60.[Pub Med][Google Scholar]
- 4.Myung DS,Jung YS,KangAJ,etal..Primary shewanella algae bacteremia mimicking Vibrio septicemia. Korean Med Sc.2009;24:1192-1194
- 5.Shah BA,Padbury JF. Neonatal sepsis: an old problem with new insights .Virulance.2014;5:170-8.[PMJ Free article][PubMed][Google Scholar]
- 6.Otsuka T,Noda T, Noguchi A,Nakamura H,Ibaraki K,Yamaoka k. Shewanella infection in decompansated liver disease:a septic case .J Gastroenterol .2007;42:87-90.[PubMed][Google scholar]

- 7.Khashe S,Janda JM.Biochemical and pathological properties of shewanella algae and shewanella putrefaciens .J Clin Microbiol.1998;36:783-7. [PMC Free article][PubMed][Google Scholar]20.
- 8.Domínguez H,Vogel BF, Gram L ,et al.. shewanella algae bacteremia in two patients with lower leg ulcers .Clin Infect Dis. 1999;22:1036-1039.
- 9.Pagani L , Lang A,Vedovelli C ,Moling O, Romantic G ,Pristera R. et al .Soft tissue infection and bacteremia caused by shewanella putrefaciens. J Clin Microbiol .2003;41:2240-1.[PMC Free article][PubMed][Google Scholar]
- 10.Brink AJ ,Van Straten A ,Van Rensburg AJ . Shewanella (Pseudomonas) putrefaciens bacteremia. Clin Infect Dis.1995;20:1327-32.[PubMed][Google Scholar]
- 11.Chen YS, Liu YC ,Yen MY , Wang JH , Wann SR ,Cheng DL .Skin and soft tissue manifestations of shewanella putrefaciens infection . Clin Infect Dis.1997;25:225-9.[PubMed][Google Scholar]
- 12.Sharma KK ,Kalawat U.Emerging infections: Shewanella – a series of five cases .J Lab Physicians.2010;2: 61-5.[PMC Free article] [PubMed][Google scholar]
- 13.Liu PY ,Lin CF,Tung KC, Wu MJ,Liu JW .et al .Clinical and microbiological features of Shewanella bacteremia in patients with hepatobiliary disease. Intern Med .2013;52:431-8.[PubMed][Google scholar]
14. VignierN ,Barreau M,Olive C ,Baubion E ,Theodose R, Hochedez P.et al. Human infection with shewanella putrefaciens and S.algae :report of 16 cases of Martinique and review of the literature. Am J Trop Med Hyg.2013;89:151-6.[PMC Free article][PubMed][Google Scholar]
15. Charles MVP, Srirangaraj S, Kali A.Australas.Neonatal sepsis caused by shewanella algae .A case report. AMJ .2015;8(2):64-66 .<http://dx.doi.org/10.4066/AMJ.2015.2292>
- 16.García -Fragoso, Lourdes MD ;García -García ,Inés MD ;Rivera, Amarilis MD. Shewanella algae Bacteremia in a preterm newborn .the pediatric Infectious Disease Journal :January2012- volume 31-Issue 1-p 104-105.

- 17.Sardelic S ,Karanovic J ,Rubic Z, et al .Late ventriculoperitoneal shunt infection caused by shewanella algae. *Pediatr Infect Dis J* .2010;29:475-477.
- 18.Iwata M ,Tateda R ,Matsumoto T ,et al ..Primary shewanella algae septicemia in patient on hemodialysis .*J Clin Microbiol*. 1999;37:2101-2105.
- 19.Tan CK ,Lai CC,Kuar WK ,et al ..Purulent pericarditis with greenish pericardial effusion caused by shewanella algae. *J Clin Microbiol* .2008;46:2817-2819.
- 20.Krsnik I ,Arribalzaga K ,Romany K J. Shewanella algae bacteremia and associated cellulites in a patient with multiple myeloma .*Haematologia (Budap)*.2002;33:79-80.
- 21.Shimizu T ,Matsumura Y.A case of bacteremia and suppurative vertebral osteomyelitis/discitis due to shewanella algae occurring after raw –fish consumption .*Kansenshogaku Zasshi*.2009;83:553-556.
- 22.Oh HS,Kum KA, Kim EC,etal..Outbreak of shewanella algae and shewanella putrefaciens infections caused by a shared measuring cup in general surgery unit in Korea .*Infect control Hosp Epidemiol*.2008;29:742-748.
- 23.Kim DM,Kang CI, Lee CS, etal .. Treatment Failure due to emergence of resistance to carbapenem during therapy for shewanella algae bacteremia .*J Clin Microbiol*.2006;44:1172-1174.